

Europäisches Patentamt
European Patent Office
Office européen des brevets



(11) **EP 1 016 710 A2**

(12) **EUROPEAN PATENT APPLICATION**

(43) Date of publication:
05.07.2000 Bulletin 2000/27

(51) Int. Cl.⁷: **C12N 1/20, C12N 15/11,
C07K 14/245**

(21) Application number: 99125263.6

(22) Date of filing: 17.12.1999

(84) Designated Contracting States:
**AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE**
Designated Extension States:
AL LT LV MK RO SI

- Aleshin, Vladimir Veniaminovich
Borovsk, 249010 (RU)
- Troshin, Petr Vladimirovich
Moscow, 115561 (RU)
- Tokhmakova, Irina Lyvovna
143513 Moscow region (RU)

(30) Priority: 30.12.1998 RU 98124016
09.03.1999 RU 99104431

(74) Representative:
**Strehl Schübel-Hopf & Partner
Maximilianstrasse 54
80538 München (DE)**

(72) Inventors:
• Livshits, Vitaliy Arkadievich
Moscow, 113208 (RU)
• Zakataeva, Natalia Pavlovna
Moscow, 117421 (RU)
• Nakanishi, Kazuo
Sakae-ku, Yokohama-shi, 247-0014 (JP)

Remarks:

The biological material has been deposited with
VKPM under numbers B-7728 / B-7729 / B-7731
/ B-7730 / B-7719 / B-7722 / B-7707 / B-7712 / B-
7708 / B-7753 / B-7754 / B-7752 / B-7713 / B-7714 /
B7748 / B-7715 / B-7716 / B-7718.

(54) **Method for producing L-amino acids**

(57) A bacterium belonging to the genus *Escherichia* and having an ability to produce an L-amino acid, wherein the ability to produce the L-amino acid is increased by increasing an expression amount of an L-amino acid excretion protein, and a method for producing the L-amino acid using the bacterium.

EP 1 016 710 A2

Description**Technical Field**

5 [0001] The present invention relates to a method for producing an amino acid. In particular, the present invention relates to an L-amino acid-producing bacterium belonging to the genus *Escherichia* and a method for producing L-amino acids, more specifically, L-glutamic acid, L-lysine, L-threonine, L-alanine, L-histidine, L-proline, L-arginine, L-valine, and L-isoleucine, using the bacterium.

10 **Background Art**

[0002] For production of an L-amino acid by fermentation, a strain isolated from the natural world or an artificial mutant of the strain has been used to improve productivity. For example, in the case of L-lysine, many artificial mutants producing L-lysine are known, and most of them are mutants resistant to S-2-aminoethylcysteine (AEC) and belong to the genus *Brevibacterium*, *Corynebacterium*, *Bacillus* or *Escherichia*. Also, there have been proposed various technics for increasing amino acid production such as use of a transformant obtained by using a recombinant DNA (U.S. Patent No. 4,278,765).

15 [0003] The technics are mostly based on enhancement of an activity of an enzyme involved in an amino acid biosynthetic pathway, conversion of the enzyme to that desensitized in inhibition and the like (As to bacterium belonging to the genus *Escherichia*, see Japanese Patent Application Laid-Open No. 56-18596 (1981) and International Publication No. WO 95/16042).

[0004] On the other hand, as an example of improvement of amino acid productivity by enhancing an amino acid excretion protein, a bacterium belonging to the genus *Corynebacterium* in which an L-lysine excretion gene, *lysE* is enhanced is known. However, as to bacteria belonging to the genus *Escherichia*, it is unknown even whether an L-amino acid excretion protein is present or not. Therefore, it is unknown whether enhancement of the L-amino acid excretion protein is effective in L-amino acid production using a bacterium belonging to the genus *Escherichia* or not.

25 [0005] Although the entire nucleotide sequence of *E. coli* strain K-12 belonging to the genus *Escherichia* has been already determined (Science, 277, 1453-1474(1997)), there are a large number of proteins of which functions are unknown.

30 **Disclosure of the Invention**

[0006] An object of the present invention is to obtain a protein participating in excretion of an L-amino acid, thereby providing a strain improved in L-amino acid productivity and an improved method for producing an L-amino acid by fermentation.

35 [0007] The inventors have conducted screening for the protein participating in excretion of an L-amino acid. As a result, the present inventors have found that a yield of an L-amino acid based on consumed sugar is increased when a particular gene is enhanced. On the basis of the finding, the present invention has been completed.

40 [0008] Thus, the present invention provides a bacterium belonging to the genus *Escherichia* and having an ability to produce an L-amino acid, wherein the ability to produce the L-amino acid is increased by increasing an expression amount of at least one protein selected from the group consisting of the following proteins of (A) to (H) (hereinafter also referred to as "the bacterium of the present invention"):

- (A) a protein having an amino acid sequence shown in SEQ ID NO: 10 in Sequence Listing;
- 45 (B) a protein which has an amino acid sequence including deletion, substitution, insertion, addition or inversion of one or several amino acids in the amino acid sequence shown in SEQ ID NO: 10 in Sequence Listing, and which has an activity of increasing the ability to produce the L-amino acid of the bacterium having the protein;
- (C) a protein having an amino acid sequence shown in SEQ ID NO: 12 in Sequence Listing;
- (D) a protein which has an amino acid sequence including deletion, substitution, insertion, addition or inversion of one or several amino acids in the amino acid sequence shown in SEQ ID NO: 12 in Sequence Listing, and which has an activity of increasing the ability to produce the L-amino acid of the bacterium having the protein;
- 50 (E) a protein having an amino acid sequence shown in SEQ ID NO: 14 in Sequence Listing;
- (F) a protein which has an amino acid sequence including deletion, substitution, insertion, addition or inversion of one or several amino acids in the amino acid sequence shown in SEQ ID NO: 14 in Sequence Listing, and which has an activity of increasing the ability to produce the L-amino acid of the bacterium having the protein;
- 55 (G) a protein having an amino acid sequence shown in SEQ ID NO: 16 in Sequence Listing; or
- (H) a protein which has an amino acid sequence including deletion, substitution, insertion, addition or inversion of one or several amino acids in the amino acid sequence shown in SEQ ID NO: 16 in Sequence Listing, and which

has an activity of increasing the ability to produce the L-amino acid of the bacterium having the protein.

[0009] The bacterium of the present invention preferably an L-lysine-producing bacterium in which an expression amount of at least one protein selected from the group consisting of the proteins (A) to (D), (G) and (H) is increased; an L-glutamic acid-producing bacterium in which an expression amount of at least one protein selected from the group consisting of the proteins (A) to (H) is increased; an L-alanine-producing bacterium in which an expression amount of at least one protein selected from the group consisting of the proteins (C) and (D) is increased; an L-valine-producing bacterium in which an expression amount of at least one protein selected from the group consisting of the proteins (C) and (D) is increased; an L-histidine-producing bacterium in which an expression amount of at least one protein selected from the group consisting of said proteins (C) to (F) is increased; an L-proline-producing bacterium in which an expression amount of at least one protein selected from the group consisting of said proteins (A) to (F) is increased; an L-threonine-producing bacterium in which an expression amount of at least one protein selected from the group consisting of said proteins (E) and (F) is increased; an L-arginine-producing bacterium in which an expression amount of at least one protein selected from the group consisting of said proteins (G) and (H) is increased; or an L-isoleucine-producing bacterium in which an expression amount of at least one protein selected from the group consisting of said proteins (C) and (D) is increased.

[0010] Preferably, in the bacterium of the present invention, a copy number of a DNA coding for said protein in a cell is increased. The DNA is preferably carried on a multicopy vector in the cell or on a transposon in the cell.

[0011] The present invention also provides a method for producing an L-amino acid, comprising the steps of:

cultivating the bacterium of the present invention in a culture medium, to produce and accumulate the L-amino acid in the medium, and
recovering the L-amino acid from the medium (hereinafter also referred to as "the bacterium of the present invention").

[0012] The method of the present invention preferably an L-lysine production method using an L-lysine-producing bacterium in which an expression amount of at least one protein selected from the group consisting of the proteins (A) to (D), (G) and (H) is increased; an L-glutamic acid production method using an L-glutamic acid-producing bacterium in which an expression amount of at least one protein selected from the group consisting of the proteins (A) to (H) is increased; an L-alanine production method using an L-alanine-producing bacterium in which an expression amount of at least one protein selected from the group consisting of the proteins (C) and (D) is increased; an L-valine production method using an L-valine-producing bacterium in which an expression amount of at least one protein selected from the group consisting of the proteins (C) and (D) is increased; an L-histidine production method using an L-histidine-producing bacterium in which an expression amount of at least one protein selected from the group consisting of said proteins (C) to (F) is increased; an L-proline production method using an L-proline-producing bacterium in which an expression amount of at least one protein selected from the group consisting of said proteins (A) to (F) is increased; an L-threonine production method using an L-threonine-producing bacterium in which an expression amount of at least one protein selected from the group consisting of said proteins (E) and (F) is increased; an L-arginine production method using an L-arginine-producing bacterium in which an expression amount of at least one protein selected from the group consisting of said proteins (G) and (H) is increased; or an L-isoleucine production method using an L-isoleucine-producing bacterium in which an expression amount of at least one protein selected from the group consisting of said proteins (C) and (D) is increased.

[0013] Preferably, in the method of the present invention, a copy number of a DNA coding for said protein in a cell of the bacterium is increased. The DNA is preferably carried on a multicopy vector in the cell, or on a transposon in the cell.

[0014] According to the present invention, an ability to produce an L-amino acid of a bacterium belonging to the genus *Escherichia* can be increased. Also, a method for producing an L-amino acid can be improved in a production rate of an L-amino acid.

[0015] The present invention will be explained in detail below. Hereinafter, an amino acid is of L-configuration unless otherwise noted.

(1) Bacterium of the present invention

[0016] The bacterium of the present invention is a bacterium belonging to the genus *Escherichia* and having an ability to produce an amino acid, in which the ability to produce the amino acid is increased by increasing an expression amount of a protein which has an activity of increasing the ability to produce the amino acid of the bacterium, or an activity of increasing resistance to an amino acid or amino acid analogue. Hereinafter, the protein is referred to as "amino acid excretion protein" for the sake of convenience. However, the term does not mean that function of the protein

is limited to amino acid excretion.

[0017] Examples of the amino acid excretion protein include a protein having an amino acid sequence shown in SEQ ID NO: 10, a protein having an amino acid sequence shown in SEQ ID NO: 12, a protein having an amino acid sequence shown in SEQ ID NO: 14 and a protein having an amino acid sequence shown in SEQ ID NO: 16.

5 [0018] The amino acid excretion protein may have selectivity to amino acid. An amino acid excretion protein appropriate for each amino acid can be determined by allowing the amino acid excretion protein to be expressed in a bacterium belonging to the genus *Escherichia* and having an ability to produce the amino acid, and measuring an increase of a yield of the amino acid or measuring an increase of a minimum inhibition concentration (MIC) of an amino acid or amino acid analogue.

10 [0019] For example, in the case of lysine, a protein having an amino acid sequence shown in SEQ ID NO: 10, 12 or 16 is effective; in the case of glutamic acid, a protein having an amino acid sequence shown in SEQ ID NO: 10, 12, 14 or 16 is effective; in the case of alanine, a protein having an amino acid sequence shown in SEQ ID NO: 12 is effective; in the case of valine, a protein having an amino acid sequence shown in SEQ ID NO: 12 is effective; in the case of histidine, a protein having an amino acid sequence shown in SEQ ID NO: 12 or 14; in the case of proline, a protein having an amino acid sequence shown in SEQ ID NO: 10, 12 or 14 is effective; in the case of threonine, a protein having an amino acid sequence shown in SEQ ID NO: 14 is effective; in the case of arginine, a protein having an amino acid sequence shown in SEQ ID NO: 16 is effective; and in the case of isoleucine, a protein having an amino acid sequence shown in SEQ ID NO: 12 is effective.

20 [0020] The term "an expression amount is increased" used herein usually means that the expression amount is larger than that in a wild strain of *E. coli* such as strain MG1655 or W3110. The term also means that when a strain is obtained by modification through genetic engineering techniques or the like, the expression amount is larger than that prior to the modification. The expression amount of the amino acid excretion protein may be determined directly by the determination of the amino acid excretion protein or indirectly by the determination of MIC of an amino acid or amino acid analogue or of amino acid productivity of a bacterium belonging to the genus *Escherichia* and having the amino acid excretion protein.

25 [0021] The method for increasing the expression amount of the amino acid excretion protein is exemplified by a method for increasing a copy number of DNA encoding the amino acid excretion protein in a cell of the bacterium.

30 [0022] For increasing the copy number in the cell, a DNA fragment coding for the amino acid excretion protein may be ligated to a vector which functions in a bacterium belonging to the genus *Escherichia* to produce a recombinant DNA, which is introduced to a host to transform it. The copy number of the gene coding for the amino acid excretion protein (amino acid excretion protein gene) in the cell of the transformant strain increases, thereby increasing the expression amount of the amino acid excretion protein. The vector is preferably a multicopy vector.

35 [0023] The increase of the copy number in the cell can be achieved by allowing plural copies of the amino acid excretion protein gene to exist on chromosomal DNA of the host. The introduction of plural copies of the amino acid excretion protein gene to chromosomal DNA of a bacterium belonging to the genus *Escherichia*, may be conducted through homologous recombination by using a sequence of which plural copies exist on the chromosomal DNA, as a target. As the sequence of which plural copies exist on the chromosomal DNA, a repetitive DNA and an inverted repeat present in a terminal portion of a transposable element may be used. Alternatively, as disclosed in Japanese Patent Application Laid-Open No. 2-109985 (1990), the plural copies can be introduced to the chromosomal DNA by making the amino acid excretion protein gene carried on a transposon and allowing the transposon to be transposed, which is preferred. According to any of the above-mentioned methods, the copy number of the amino acid excretion protein gene in the transformant strain increases, thereby increasing the expression amount of the amino acid excretion protein.

40 [0024] The multicopy vector is exemplified by plasmid vectors such as pBR322, pMW118, pUC19 or the like, and phage vectors such as λ 1059, λ BF101, M13mp9 or the like. The transposon is exemplified by Mu, Tn10, Tn5 or the like.

45 [0025] The introduction of a DNA into a bacterium belonging to the genus *Escherichia* can be performed, for example, by a method of D. M. Morrison (Methods in Enzymology 68, 326 (1979)) or a method in which recipient bacterial cells are treated with calcium chloride to increase permeability of DNA (Mandel, M. and Higa, A., J. Mol. Biol., 53, 159 (1970)) and the like.

50 [0026] Besides the above-mentioned gene amplification, the increase of the expression amount of the amino acid excretion protein can be also achieved by replacing an expression regulatory sequence such as a promoter of the amino acid excretion protein gene with stronger one (see Japanese Patent Application Laid-Open No. 1-215280 (1989)). For example, *lac* promoter, *trp* promoter, *tac* promoter, P_R promoter and P_L promoter of lambda phage, and the like are known as a strong promoter. The replacement with the promoter enhances expression of the amino acid excretion protein, thereby increasing the expression amount of the amino acid excretion protein. The enhancement of the expression regulatory sequence may be combined with the increase of the copy number of the amino acid excretion protein.

55 [0027] In the bacterium of the present invention, expression amounts of plural amino acid excretion proteins may be increased.

[0028] The amino acid excretion protein is encoded by genes which are known as *yahN* gene, *yeaS* gene, *yfiK* gene and *yggA* gene and of which functions are unknown. Therefore, the DNA encoding the amino acid excretion protein can be obtained by synthesizing primers based on the known sequences (for example, the entire nucleotide sequence of chromosome of *Escherichia coli* strain K-12 has been already determined (Science, 277, 1453-1474(1997))), and conducting amplification by PCR using chromosomal DNA of a bacterium belonging to the genus *Escherichia* as a template. Also, the object DNA fragment can be selected by hybridization from a chromosomal DNA library of a bacterium belonging to the genus *Escherichia* by preparing a probe based on the known sequences. Alternatively, the DNA encoding the amino acid excretion protein may be synthesized based on the known sequences. The nucleotide sequence of the DNA encoding the amino acid excretion protein is exemplified by that shown in SEQ ID NO: 9, 11, 13 or 15 in the Sequence Listing.

[0029] Methods for preparation of chromosomal DNA, preparation of chromosomal DNA library, hybridization, PCR, preparation of plasmid DNA, digestion and ligation of DNA, transformation, selection of an oligonucleotide as a primer and the like may be ordinary methods well known to one skilled in the art. These methods are described in Sambrook, J., Fritsch, E. F., and Maniatis, T., "Molecular Cloning A Laboratory Manual, Second Edition", Cold Spring Harbor Laboratory Press (1989) and the like.

[0030] The amino acid excretion protein may comprise substitution, deletion, insertion, addition or inversion of one or several amino acids at one or a plurality of positions, provided that the activity of increasing the ability to produce the amino acid of the bacterium belonging to the genus *Escherichia* and having the protein is not deteriorated. The term "several" may vary depending on a position in a steric structure of the protein and a kind of an amino acid residue. It is because some amino acids such as isoleucine and valine have high similarity to each other, and a difference between such the amino acids does not largely affect the steric structure of the protein.

[0031] The DNA which codes for the substantially same protein as the amino acid excretion protein as described above, may be obtained, for example, by modifying the nucleotide sequence, for example, by means of the site-directed mutagenesis method so that one or more amino acid residues at a specified site involve substitution, deletion, insertion, addition or inversion. The DNA modified as described above may be obtained by the conventionally known mutation treatment. The mutation treatment includes a method for treating a DNA coding for the amino acid excretion protein *in vitro*, for example, with hydroxylamine, and a method for treating a microorganism, for example, a bacterium belonging to the genus *Escherichia*, harboring a DNA coding for the amino acid excretion protein with ultraviolet irradiation or a mutating agent such as N-methyl-N'-nitro-N-nitrosoguanidine (NG) and nitrous acid usually used for the mutation treatment.

[0032] The substitution, deletion, insertion, addition or inversion of the one or more amino acid residues includes a naturally-occurring mutation or variation which is resulted from a difference between individual microorganisms having the amino acid excretion protein and a difference between species, strains or the like.

[0033] The DNA, which codes for substantially the same protein as the amino acid excretion protein, can be obtained by allowing a DNA having the mutation as described above to be expressed in a cell of an appropriate bacterium belonging to the genus *Escherichia*, and investigating the increase of amino acid productivity of the cell.

[0034] Also, the DNA, which codes for substantially the same protein as the amino acid excretion protein, can be obtained by isolating a DNA which hybridizes with DNA having, for example, a nucleotide sequence shown in SEQ ID NO: 9, 11, 13 or 15 in Sequence Listing under stringent conditions, and which codes for a protein having the activity of increasing the ability to produce the amino acid of the bacterium belonging to the genus *Escherichia*, from DNAs encoding the amino acid excretion proteins having mutations or cells containing the DNAs. The term "stringent conditions" referred to herein means a condition under which a specific hybrid is formed, and a non-specific hybrid is not formed. It is difficult to clearly express this condition by using any numerical value. However, for example, the stringent conditions include a condition under which DNAs having high homology, for example, DNAs having homology of not less than 70% with each other are hybridized, and DNAs having homology lower than the above with each other are not hybridized, or a condition of a salt concentration corresponding to 60°C, 1x SSC, 0.1% SDS, preferably 0.1x SSC, 0.1% SDS which is a washing condition of ordinary Southern hybridization.

[0035] Although there may be a gene in which a stop codon is made in the middle, or a gene encoding a protein losing the activity due to mutation of the active center among the genes which hybridize under such the condition, such genes can be easily eliminated by ligating the genes to a commercially available activity-expression vector and determining the activity of increasing the ability to produce the amino acid of the bacterium belonging to the genus *Escherichia* as described above.

[0036] The term "DNA coding for a protein" used herein means a DNA of which one of strands codes for the protein when the DNA is double-stranded.

[0037] By increasing an expression amount of an amino acid excretion protein in an amino acid-producing bacterium belonging to the genus *Escherichia* as described above, a produced amount of the amino acid can be increased. As the bacterium belonging to the genus *Escherichia* in which the expression amount of the amino acid excretion protein is to be increased, strains which have abilities to produce desired amino acids (amino acid productivities) are used.

Besides, an ability to produce an amino acid may be imparted to a bacterium in which the expression amount of the amino acid excretion protein is increased. Examples of amino acid-producing bacteria belonging to the genus *Escherichia* include *E. coli* AJ13199 (FR patent No. 2747689), and those obtainable from known materials (e.g., *E. coli* W3110 (tyrA)/pCABD2, *E. coli* VL614, *E. coli* VL2054, *E. coli* VL2160, *E. coli* VL2151, *E. coli* W3350 argE::Tn10/pKA10 as described in the Examples below).

[0038] For reference, the amino acid excretion protein according to the present invention was identified for the first time as described below.

[0039] The present inventors have identified *rhtB* and *rhtC* as threonine excretion protein genes of a bacterium belonging to the genus *Escherichia*. The present inventors searched databases based on a hypothesis that amino acid excretion proteins may share a common structure. Namely, BLAST and PSI-BLAST search (Altschul, S.F. et al., *Nucleic Acids Res.*, 25, 3389-3402(1997)) for homology of a protein encoded by *rhtB* was performed in GenBank CDS, PDB, SWISS-PROT, Spupdate and PIR. Tblastn search was performed in unfinished microbial genomes. BLITZ search (Sturrock, S.S., and Collins, J.F., Mpsch version 1.3. Biocomputing research unit University of Edinburgh, UK (1993)) was performed in SWALL database. SMART search (Ogiwara, I. et al., *Protein Sci.*, 5, 1991-1999 (1996)) was performed in the databases of translations and SWISS-PROT. From the samples of more than 60 sequences found, YeaS (corresponding to f212 of ACCESSION No. AE000274 in GenBank), YahN (corresponding to f223 of ACCESSION No. AE000140 in GenBank), YfiK (corresponding to o195 of ACCESSION No. AE000344 in GenBank) and YggA (corresponding to f211 of ACCESSION No. AE000375 in GenBank) remained as proteins which may have similar function to *RhtB*, among those originating from *E. coli*. Since functions of any of these genes were unknown, the genes were actually obtained, and effects thereof on MIC of amino acids and amino acid analogues and on amino acid production were examined by enhancing activities thereof. As a result, an effect of increasing MIC of some amino acids and analogues was found with respect to YeaS, YfiK, YahN and YggA. Further examination has revealed that proteins encoded by these genes exhibit an effect of increasing an amino acid accumulation, although they may have some amino acid selectivities.

(2) Method of the present invention

[0040] The method of the present invention comprises the steps of cultivating the bacterium of the present invention, in a culture medium, to produce and accumulate the amino acid in the medium, and recovering the amino acid from the medium.

[0041] Suitable amino acids include lysine, glutamic acid, alanine, valine, homoserine, proline, and threonine.

[0042] In the method of present invention, the cultivation of the bacterium belonging to the genus *Escherichia*, the collection and purification of amino acid from the liquid medium may be performed in a manner similar to those of the conventional method for producing an amino acid by fermentation using a bacterium. A medium used in cultivation may be either a synthetic medium or a natural medium, so long as the medium includes a carbon and a nitrogen source and minerals and, if necessary, nutrients which the bacterium used requires for growth in appropriate amounts. The carbon source may include various carbohydrates such as glucose and sucrose, and various organic acids. Depending on assimilatory ability of the used bacterium, alcohol including ethanol and glycerol may be used. As the nitrogen source, ammonia, various ammonium salts such as ammonium sulfate, other nitrogen compounds such as amines, a natural nitrogen source such as peptone, soybean hydrolyte and digested fermentative microbe are used. As minerals, monopotassium phosphate, magnesium sulfate, sodium chloride, ferrous sulfate, manganese sulfate, calcium carbonate are used.

[0043] The cultivation is preferably culture under an aerobic condition such as a shaking culture, and an aeration and stirring culture. The temperature of culture is usually 20 to 40°C, preferably 30 to 38°C. The pH of the culture is usually between 5 and 9, preferably between 6.5 and 7.2. The pH of the culture can be adjusted with ammonia, calcium carbonate, various acids, various bases, and buffers. Usually, a 1 to 3-day cultivation leads to the accumulation of the target amino acid in the medium.

[0044] Recovering the amino acid can be performed by removing solids such as cells from the medium by centrifugation or membrane filtration after cultivation, and then collecting and purifying the target amino acid by ion exchange, concentration and crystalline fraction methods and the like.

Best Mode for Carrying Out the Invention

[0045] The present invention will be more concretely explained below with reference to Examples.

Example 1. Preparation of the DNA fragments which code for amino acid excretion proteins.

[0046] The entire nucleotide sequence of chromosome of *E. coli* strain K-12 has been determined (Science, 277,

1453-1474, 1997). Based on the reported nucleotide sequence, primers were synthesized and the genes *yahN*, *yfiK*, *yeaS* and *yggA* were amplified by PCR.

(1). Chromosomal DNA of the *E. coli* strain MG1655 was used as a template.

[0047] The chromosomal DNA was prepared by an ordinary method (Sambrook, J., Fritsch E. F. and Maniatis T. (1989) Molecular cloning: a laboratory manual, 2nd ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N. Y.). In the PCR reaction, a standard condition described in "PCR protocols. Current methods and applications". (White, B.A., ed. Humana Press, Totowa, New Jersey, 1993) was used. The obtained PCR products were purified by an ordinary method and digested with restriction enzymes as described below.

[0048] The *yahN* gene was amplified by using the primers No.1 and No. 2.

Primer No.1: gtgtggaaccgacgccggat (a sequence complementary to a sequence of from 1885 base to 1904 base in a nucleotide sequence registered under ACCESSION No. AE000140 in GenBank; SEQ ID NO: 17), and

Primer No.2: tgtgtatggtacggggtcgag (a sequence of from 223 base to 245 base in the same; SEQ ID NO: 18).

[0049] The obtained PCR product after purification was digested with restriction enzymes *Pst*I and *Stu*I and ligated to vector pUC21 (Vieira, Messing, Gene, 100, 189-194, 1991) digested with the enzymes *Pst*I and *Eco*RV by using a ligation kit. Then, transformation of competent cells of *E. coli* TG1 (Sambrook, J., Fritsch E. F. and Maniatis T. (1989) Molecular cloning: a laboratory manual, 2nd ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N. Y.) with the product was conducted and the cells were spread on L medium (10 g/l Bacto trypton, 5 g/l Yeast extract, 5 g/l NaCl, 15 g/l agar, pH 7.0) containing 10 mg/ml IPTG (isopropyl- β -D-thiogalactopyranoside) and 40 mg/ml X-gal (5-bromo-4-chloro-3-indolyl- β -D-galactoside) and 100 mg/ml ampicillin, and cultured overnight. Appeared white colonies were picked up and subjected to single colony isolation to obtain transformants. Plasmid was prepared from the transformants using an alkali extraction method and designated as pYAHN.

[0050] The *yeaS* gene was amplified by using the primers No.3 and No. 4.

Primer No.3: ctttgcaatcccgtctccc (a sequence complementary to a sequence of from 7683 base to 7702 base in a nucleotide sequence registered under ACCESSION No AE000274 in GenBank; SEQ ID NO: 19);

Primer No.4: gccccatgcataacggaaag (a sequence of from 5542 base to 5561 base in the same; SEQ ID NO: 19).

[0051] The obtained PCR product after purification was digested with a restriction enzyme *Ava*I and ligated to vector pUC19. After transformation of *E. coli* TG1 as above, the plasmid designated as pYEAS was obtained.

[0052] The *yfiK* gene was amplified by using the primers No.5 and No.6.

Primer No.5: gaagatctttaggcgggataaggcg (a sequence of from 4155 base to 4177 base in a nucleotide sequence registered under ACCESSION No AE000344 in GenBank, with a restriction enzyme *Bgl*II site added at the 5'-end thereof; SEQ ID NO: 21)

Primer No.6: tggttttaccaattggccgc (a sequence complementary to a sequence of from 6307 base to 6326 base in the same; SEQ ID NO: 22).

[0053] The obtained PCR product after purification was digested with restriction enzymes *Bgl*II and *Mun*I and ligated to vector pUC21 digested with restriction enzymes *Bgl*II and *Eco*RI. After transformation of *E. coli* TG1 as above, the plasmid designated pYFIK was obtained.

[0054] The *yggA* gene was amplified by using the primers No.7 and No.8.

Primer No.7: acttctccgcgagccagttc (a sequence complementary to a sequence of from 9606 base to 9626 base in a nucleotide sequence registered under ACCESSION No AE000375 in GenBank; SEQ ID NO: 23).

Primer No.8: ggcaagcttagcgctctgtt (a sequence of from 8478 base to 8498 base in the same; SEQ ID NO: 24).

[0055] The obtained PCR product after purification was digested with restriction enzymes *Hind*III and *Cla*I and ligated to vector pOK12 (Vieira, Messing, Gene, 100, 189-194, 1991) digested with the same restriction enzymes. After transformation of *E. coli* TG1 as above, the plasmid designated pYGGA was obtained.

(2). Chromosomal DNA of the *E. coli* strain W3110 was used as a template.

[0056] The *yahN* gene was amplified by using the primers No.9 (SEQ ID NO 1) and No. 10 (SEQ ID NO.2)
 [0057] The *yeaS* gene was amplified by using the primers No.11 (SEQ ID NO 3) and No.12 (SEQ ID NO 4)
 [0058] The *yfiK* gene was amplified by using the primers No.13 (SEQ ID NO 5) and No.14 (SEQ ID NO 6).
 [0059] The *yggA* gene was amplified by using the primers No.15 (SEQ ID NO 7) and No.16 (SEQ ID NO 8)
 [0060] The obtained PCR product was purified, digested with restriction enzymes *SacI* and *XbaI* (*EcoRI* and *PstI* for *yggA*), and ligated to plasmid pMW118 (Nippon Gene). The plasmid into which a DNA fragment of which sequence was identical to the reported sequence was inserted was designated as follows:

One carrying *yahN*: pMW118::*yahN*
 One carrying *yeaS*: pMW118::*yeaS*
 One carrying *yfiK*: pMW118::*yfiK*
 One carrying *yggA*: pMW118::*yggA*

Example 2. Effect of the *yahN*, *yeaS*, *yfiK*, and *yggA* DNA fragments amplification on the *E. coli* TG1 resistance to some amino acids and amino acid analogues.

[0061] The homology of the *yeaS*, *yfiK*, *yahN* and *yggA* gene products with the lysine transporter, LysE, of *Corynebacterium glutamicum* (Vrljic et al., Mol. Microbiol., 22, 815-826, 1996) and RhtB protein involved in homoserine excretion, indicates the analogues function for these proteins. It is well known that the increased expression of the genes involved in antibiotic and heavy metal efflux increases the level of resistance to the drugs (Nikaido, H. J. Bacteriology, 178, 5853-5859, 1996). Therefore, the effect of the pYEAS, pYAHN, pYFIK, and pYGGA plasmids on susceptibility of the strain TG1 to some amino acids and amino acid analogues was tested. Overnight cultures of the *E. coli* strains TG1/pYEAS, TG1/pYAHN, TG1/pYFIK, TG1/pYGGA and of the control strains TG1/pUC21, TG1/pUC19 and TG1/pOK12 grown in M9 minimal medium with an appropriate antibiotic on a rotary shaker (10^9 cfu/ml) were diluted 1:100 in M9 minimal medium and grown for 5 h in the same medium. Then the log phase cultures thus obtained were diluted and about 10^4 alive cells were applied to well-dried test plates with M9 agar containing doubling increments of amino acids or analogues. Thus the minimum inhibition concentration (MIC) of these compounds were examined.
 [0062] The results are shown in Table 1. It follows from the Table 1 that multiple copies of *yfiK* gene conferred increased resistance to proline, homoserine, histidine, threonine, glutamate, lysine, α -amino- β -hydroxyvaleric-acid (AHVA), S-(2-aminoethyl)-L-cysteine (AEC) and α -aminobutyric acid; multiple copies of *yahN* gene conferred increased resistance to proline, multiple copies of *yeaS* gene conferred increased resistance to threonine, homoserine, lysine, glutamate, histidine, proline and α -aminobutyric acid; multiple copies of *yggA* gene conferred increased resistance to S-(2-aminoethyl)-L-cysteine (AEC), lysine, and arginine. These results indicate that except for *YahN*, every of the presumed transporters have specificity to several substrates (amino acids and amino acid analogues), or may show non-specific effects as a result of amplification.

Table 1

| Substrate | MIC (μ g/ml) for <i>E. coli</i> TG1, harboring the plasmid | | | | |
|-----------------------|---|-------|-------|-------|-------|
| | pUC21 | pYFIK | pYAHN | pYEAS | pYGGA |
| L-homoserine | 500 | 1000 | 500 | 1000 | 500 |
| L-threonine | 30000 | 40000 | 30000 | 50000 | 30000 |
| L-lysine | 5000 | 7500 | 5000 | 7500 | 15000 |
| L-glutamate (Na salt) | 5000 | 10000 | 5000 | 20000 | 5000 |
| L-histidine | 5000 | 10000 | 5000 | 30000 | 5000 |
| L-valine | 0.5 | 0.5 | 0.5 | 0.5 | 0.5 |
| L-proline | 1000 | 5000 | 2000 | 2000 | 1000 |
| L-arginine | 10000 | 10000 | 10000 | 10000 | 20000 |
| AHVA | 100 | 200 | 100 | 100 | 100 |
| AEC | 5 | 10 | 5 | 5 | 200 |

Table 1 (continued)

| Substrate | MIC ($\mu\text{g/ml}$) for <i>E. coli</i> TG1, harboring the plasmid | | | | |
|-----------------------------|--|-------|-------|--------|-------|
| | pUC21 | pYFIK | pYAHN | pYEAS | pYGGA |
| α -aminobutyric acid | 2500 | 5000 | 2500 | >10000 | 2500 |
| 4-aza-DL-leucine | 100 | 100 | 100 | 100 | 100 |

Example 3. Effect of *yeaS*, *yahN*, and *yfiK* DNA fragments amplification on glutamic acid production.

[0063] The *E. coli* strain AJ13199 (FR patent No. 2747689) was transformed with the vector pUC21 and each of the plasmids pYAHN, pYEAS and pYFIK. Thus the strains AJ13199/pUC21 (VKPM B-7728), AJ13199/pYAHN (VKPM B-7729), AJ13199/pYEAS (VKPM B-7731), and AJ13199/pYFIK (VKPM B-7730) were obtained.

[0064] These strains were each cultivated at 37°C for 18 hours in a nutrient broth with 100 mg/l ampicillin, and 0.3 ml of the obtained culture was inoculated into 3 ml of a fermentation medium containing 100 mg/l ampicillin, in a 20 x 200 mm test tube, and cultivated at 37°C for 48 hours with a rotary shaker. After the cultivation, an accumulated amount of glutamic acid in the medium was determined by known method.

[0065] The composition of the fermentation medium (g/l):

| | |
|--|---|
| Glucose | 80 |
| $(\text{NH}_4)_2\text{SO}_4$ | 22 |
| K_2HPO_4 | 2 |
| NaCl | 0.8 |
| $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ | 0.8 |
| $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$ | 0.02 |
| $\text{MnSO}_4 \cdot 5\text{H}_2\text{O}$ | 0.02 |
| Thiamine HCl | 0.0002 |
| Yeast extract | 1.0 |
| CaCO_3 | 30.0 (dry-heat-sterilized at 180°C for 2 h) |
| (Glucose and K_2HPO_4 separately sterilized) | |

[0066] The results are shown in Table 2. As shown in Table 2, the strains AJ13199/pYAHN, AJ13199/pYEAS, and AJ13199/pYFIK accumulated glutamic acid in a larger amount than the strain AJ13199/pUC21 in which an expression amount of amino acid excretion proteins was not enhanced.

Table 2

| Strain | Glutamic acid, g/l |
|---------------|--------------------|
| AJ13199/pUC21 | 21.9 |
| AJ13199/pYAHN | 27.9 |
| AJ13199/pYEAS | 29.7 |
| AJ13199/pYFIK | 28.4 |

Example 4. Effect of *yeaS*, *yahN*, and *yfiK* DNA fragments amplification on lysine production.

[0067]

(1). As the lysine-producing bacterium belonging to the genus *Escherichia*, *E. coli* strain W3110 (TyrA) described

in European Patent Publication No. 488424 to which plasmid pCABD2 was introduced, described in International Publication No. WO 95/16042) was used. Specifically, plasmid pCABD2, and each of the plasmid pMW118::yahn, pMW118::yeaS, pMW118::yfiK and pMW118 were introduced to *E. coli* strain W3110 (TyrA) to obtain the following strains:

W3110 (tyrA)/pCABD2+pMW118::yahn
W3110 (tyrA)/pCABD2+pMW118::yeaS
W3110 (tyrA)/pCABD2+pMW118::yfiK
W3110 (tyrA)/pCABD2+pMW118.

Lysine productivity of these strains was estimated by culture. The composition of the used medium was as follows (g/l):

| | |
|---|------|
| Glucose | 40.0 |
| MgSO ₄ · 7H ₂ O | 1.0 |
| (NH ₄) ₂ SO ₄ | 16.0 |
| K ₂ HPO ₄ | 1.0 |
| FeSO ₄ · 7H ₂ O | 0.01 |
| MnSO ₄ · 7H ₂ O | 0.01 |
| Yeast extract (Difco) | 2.0 |
| Tyrosine | 0.1 |
| Adjusted to pH 7.0 and autoclaved at 115° C for 10 minutes. (Glucose and MgSO ₄ · 7H ₂ O separately sterilized) | |
| Pharmacoepial CaCO ₃ 25 g/l (dry-heat-sterilized at 180°C for 2 h) | |

As antibiotics, 20 mg/l of streptomycin and 50 mg/l of ampicillin were added depending on a kind of a plasmid. Cultivation was conducted at 37°C for 30 hours with agitation at 115 rpm. The results are shown in Table 3.

Table 3

| Strain | Lysine, g/l | Yield, (%) |
|-----------------------------------|-------------|------------|
| W3110(tyrA) | 0.08 | 0.2 |
| W3110(tyrA)/pCABD2 + pMW118 | 12.2 | 30.5 |
| W3110(tyrA)/pCABD2 + pMW118::yahn | 13.8 | 34.5 |
| W3110(tyrA)/pCABD2 + pMW118::yeaS | 12.7 | 31.8 |
| W3110(tyrA)/pCABD2 + pMW118::yfiK | 12.2 | 30.5 |

The result in Table 3 shows that the produced amount and the yield based on consumed sugar of lysine is increased by enhancement of YahN and YeaS.

(2). As the lysine-producing bacterium belonging to the genus *Escherichia*, *E. coli* strain VL614 was used. This strain is a derivative of the known *E. coli* strain VL613 (SU Patent No. 1354458). In turn, the strain VL613 was obtained from the known strain Gif102 (Theze, J. and Saint Girons. J.Bacteriol., 118, 990-998, 1974) in the three steps:

At the first step the mutants resistant to 2 mg/ml S-(2-aminoethyl)-L-cysteine were selected and among them the strain VL611 was found capable to produce L-lysine.

At the second step the genes involved in sucrose utilization and located on the transposon Tn2555 (Doroshenko et al., Mol. Biologiya, 22, 645-658, 1988), were introduced into VL611 using phage P1-mediated trans-

duction giving the strain VL612.

At the third step, the mutation *rhtA23* from the strain VKPM B-3996, conferring resistance to threonine and homoserine (US Patent No. 5,175,107) was introduced into VL612 by phage P1 transduction giving the strain VL613.

5

[0068] The *E. coli* strain VL614 was obtained by transduction of the wild-type allele of the *rhtA* gene from the *E. coli* strain VKPM B-6204 (MG1655 *zb3058::Tn10*) to VL613. Transductants were selected on L-medium containing 10 mg/l tetracyclin, and among them the strain VL614 (*rhtA*⁺) sensitive to 10 g/l homoserine was found.

10

[0069] The strain VL614 was transformed with the pYGGGA plasmid or with the pOK12 vector to obtain strains VL614/pYGGGA (VKPM B-7719) and VL614/pOK12 (VKPM B-7722).

15

[0070] These strains were each cultivated at 37°C for 18 hours in a nutrient broth with 50 mg/l kanamycin, and 0.3 ml of the obtained culture was inoculated into 3 ml of a fermentation medium (Example 3) containing 0.3 g/l threonine, 0.3 g/l methionine and 50 mg/l kanamycin, in a 20 x 200 mm test tube, and cultivated at 37°C for 48 hours with a rotary shaker. After the cultivation, each accumulated amount of lysine and glutamate in the medium was determined by the known method.

[0071] The results are shown in Table 4.

Table 4

20

| Strain | Lysine, g/l | Glutamate, g/l |
|--------------|-------------|----------------|
| VL614/pOK12 | 2.6 | 0.8 |
| VL614/pYGGGA | 3.6 | 2.2 |

25

[0072] As shown in Table 4, the strain VL614/pYGGGA accumulated lysine in a larger amount than the strain VL614/pOK12 in which the *yggA* gene was not enhanced. Besides, the strain VL614/pYGGGA accumulated more glutamic acid than the strain VL614/pOK12.

30

Example 5. Effect of *yeaS*, *yahN*, and *yfiK* DNA fragments amplification on threonine, alanine, valine and isoleucine production.

[0073] As the threonine-producing bacterium belonging to the genus *Escherichia*, the *E. coli* strain VL2054 was used. This strain was derived from the known *E. coli* strain VKPM B-3996 (US Patent No. 5,175,107) as follows.

35

[0074] Initially, a new recipient strain was constructed in several steps:

- The plasmidless derivative of the strain VKPM B-3996 was selected after spontaneous elimination of pVIC40 plasmid.
- The wild-type allele of the *rhtA* gene from the *E. coli* strain VKPM B-6204 (MG1655 *zb3058::Tn10*) was introduced into the thus obtained strain by phage P1 mediated transduction as in the Example 4.
- A mutation inactivating *kan* gene of the Tn5 transposon inserted into the *tdh* gene was obtained after NG mutagenesis and selection of kanamycin-sensitive cells still unable to degrade threonine. Thus the strain VL2053 was obtained.

45

[0075] On the other hand, the threonine operon from pVIC40 was cloned into integrative Mud vector under the P_R promoter of the phage lambda. In addition, the *cat* gene of Tn9 conferring the resistance to chloramphenicol was cloned into the same vector. The construct thus obtained was inserted into the chromosome of the *E. coli* strain C600 by use of the known method (US Patent No. 5,595,889) and transduced from the thus obtained strain to VL2053, giving the new plasmidless threonine-producing strain VL2054. This strain accumulated in culture broth also alanine, valine and isoleucine.

50

[0076] The strain VL2054 was transformed with each of the plasmids pYEAS, pYFIK, and with the vector pUC21 to obtain *E. coli* strains VL2054/pYEAS (VKPM B-7707), VL2054/pYFIK (VKPM B-7712) and VL2054/pUC21 (VKPM B-7708).

55

[0077] These strains were each cultivated at 37°C for 18 hours in a nutrient broth with 100 mg/l ampicillin, and 0.3 ml of the obtained culture was inoculated into 3 ml of a fermentation medium (Example 3) containing 100 mg/l ampicillin, in a 20 x 200 mm test tube, and cultivated at 37°C for 48 hours with a rotary shaker. After the cultivation, each accumulated amount of threonine, alanine, valine and isoleucine in the medium was determined by known method.

[0078] The results are shown in Table 5.

[0079] As shown in Table 5, the strain VL2054/pYFIK accumulated threonine in a larger amount than the strain VL2054/pUC21 in which the *yfiK* gene was not enhanced. Besides, the strain VL2054/pYEAS accumulated more alanine, valine and isoleucine than the strain VL2054/pUC21 in which the *yeaS* gene was not enhanced.

Table 5

| Strain | Amino acid accumulation, g/l | | | |
|--------------|------------------------------|---------|--------|------------|
| | Threonine | Alanine | Valine | Isoleucine |
| VL2054/pUC21 | 5.8 | 0.4 | 0.31 | 0.15 |
| VL2054/pYEAS | 5.2 | 1.4 | 0.52 | 0.45 |
| VL2054/pYFIK | 8.8 | 0.5 | 0.22 | 0.14 |

Example 6. Effect of *yeaS* and *yfiK* DNA fragments amplification on histidine production.

[0080] As the histidine-producing bacterium belonging to the genus *Escherichia*, the strains *E. coli* VL2160 was used. This strain was obtained on the basis of the known strain NK5526 *hisG::Tn10* (VKPM B-3384) by phage P1-mediated transduction of the *hisG^R* mutation desensitizing ATP-phosphoribosyltransferase from the strain CC46 (Astvatsaturianz et al., Genetika, 24, 1928-1934, 1988). The strain *E. coli* VL2160 was transformed with each of the plasmids pYEAS, pYFIK, and with the vectors pUC21 to obtain *E. coli* strains VL2160/pYEAS (VKPM B-7753), *E. coli* VL2160/pYFIK (VKPM B-7754), *E. coli* VL2160/pUC21 (VKPM B-7752).

[0081] These strains were each cultivated at 37°C for 18 hours in a nutrient broth with 100 mg/l ampicillin, and 0.3 ml of the obtained culture was inoculated into 3 ml of the fermentation medium (Example 3) containing an increased amount of yeast extract (3 g/l) and 100 mg/l ampicillin, in a 20 x 200 mm test tube, and cultivated at 34°C for 68 hours with a rotary shaker.

[0082] After the cultivation, an accumulated amount of histidine in the medium was determined by known method. The results are shown in Table 6.

Table 6

| Strain | Histidine, g/l |
|--------------|----------------|
| VL2160/pUC21 | 1.2 |
| VL2160/pYEAS | 1.8 |
| VL2160/pYFIK | 1.4 |

[0083] As shown in Table 6, the strains *E. coli* VL2160/pYEAS and *E. coli* VL2160/pYFIK accumulated histidine in a larger amount than the strain *E. coli* VL2160/pUC21 in which the *yeaS* and *yfiK* genes were not enhanced.

Example 7. Effect of *yahN*, *yfiK* and *yeaS* DNA fragments amplification on proline production.

[0084] As the proline-producing bacterium belonging to the genus *Escherichia*, the strain VL2151 (W3350 *proB⁺ ΔputAP Tn10*) was used. This strain was obtained by transduction into W3350 of *ΔputAP* mutation linked to Tn10 and selecting tetracycline-resistant transductants unable to utilize proline as a sole carbon source. The thus obtained strain W3350 *ΔputAP Tn10* was mutagenized with NG and mutants resistant to 20 mg/l of 3,4-dehydro-DL-proline were selected. Among them the strain VL2151 (W3350 *proB⁺ ΔputAP Tn10*) was found capable to produce proline.

[0085] The strain *E. coli* VL2151 was transformed with each of the plasmids pYEAS, pYFIK, pYAHN and with the vectors pUC21 to obtain *E. coli* strains VL2151/pYEAS (VKPM B-7714), VL2151/pYFIK (VKPM B-7713), VL2151/pYAHN (VKPM B-7748) and *E. coli* VL2151/pUC21 (VKPM B-7715).

[0086] These strains were each cultivated at 37°C for 18 hours in a nutrient broth with 100 mg/l ampicillin, and 0.3 ml of the obtained culture was inoculated into 3 ml of a fermentation medium (Example 3) containing 100 mg/l ampicillin, in a 20 x 200 mm test tube, and cultivated at 37°C for 48 hours with a rotary shaker. After the cultivation, an accumulated amount of proline in the medium was determined by known method. The results are shown in Table 7.

Table 7

| Strain | Proline, g/l |
|--------------|--------------|
| VL2151/pUC21 | 1.8 |
| VL2151/pYAHN | 2.2 |
| VL2151/pYEAS | 2.1 |
| VL2151/pYFIK | 2.5 |

[0087] As shown in Table 7, the strains *E. coli* VL2151/pYFIK, *E. coli* VL2151/pYAHN and *E. coli* VL2151/pYEAS accumulated proline in a larger amount than the strain *E. coli* VL2151/pUC21 in which the *yfiK*, *yahN* and *yeaS* genes were not enhanced. The amplification of *yfiK* gene had the most pronounced effect.

Example 8. Effect of *yggA* DNA fragments amplification on arginine production.

[0088] As arginine-producing bacterium belonging to the genus *Escherichia*, the strain W3350 *argE*::Tn10/pKA10 was used. This strain harbors a plasmid, pKA10, containing DNA region from *Corynebacterium* (*Brevibacterium*) *flavum* which complements at least *argA* and *argE* mutations in the recipient strain of *E. coli* K-12 (Kharitonov A. and Tarasov A.P. Molecular Genetics, Microbiology and Virology. No.9, 29-33, 1986).

[0089] The strain *E. coli* W3350 *argE*::Tn10/pKA10 was transformed with the plasmid pYGGA, or with the vector pOK12 to obtain the strains *E. coli* W3350 *argE*::Tn10/pKA10, pYGGA (VKPM B-7716) and *E. coli* W3350 *argE*::Tn10/pKA10, pOK12 (VKPM B-7718).

[0090] The thus obtained transformants were each cultivated at 37°C for 18 hours in a nutrient broth with 100 mg/l ampicillin and 50 mg/l kanamycin, and 0.3 ml of the obtained culture was inoculated into 3 ml of a fermentation medium (Example 3) containing 100 mg/l ampicillin and 50 mg/l kanamycin, in a 20 x 200 mm test tube, and cultivated at 37°C for 48 hours with a rotary shaker. After the cultivation, an accumulated amount of arginine in the medium was determined by known method.

[0091] The results are shown in Table 8.

Table 8

| Strain | Arginine, g/l |
|---------------------------------------|---------------|
| W3350 <i>argE</i> ::Tn10/pKA10, pOK12 | 0.11 |
| W3350 <i>argE</i> ::Tn10/pKA10, pYGGA | 0.46 |

[0092] As shown in Table 8, the strains *E. coli* W3350 *argE*::Tn10/pKA10, pYGGA accumulated arginine in a larger amount than the strain *E. coli* W3350 *argE*::Tn10/pKA10, pUC21 in which the *yggA* gene was not enhanced.

[0093] The following *E. coli* strains have been deposited (according to international deposition based on Budapest Treaty) in the Russian National Collection of Industrial Microorganisms (VKPM) on December 29, 1998 under the accession numbers shown in parenthesis.

AJ13199/pUC21 (VKPM B-7728)
 AJ13199/pYAHN (VKPM B-7729)
 AJ13199/pYEAS (VKPM B-7731)
 AJ13199/pYFIK (VKPM B-7730)
 VL614/pYGGA (VKPM B-7719)
 VL614/pOK12 (VKPM B-7722)
 VL2054/pYEAS (VKPM B-7707)
 VL2054/pYFIK (VKPM B-7712)
 VL2054/pUC21 (VKPM B-7708)
 VL2160/pYEAS (VKPM B-7753)
 VL2160/pYFIK (VKPM B-7754)
 VL2160/pUC21 (VKPM B-7752)

EP 1 016 710 A2

VL2151/pYFIK (VKPM B-7713)

VL2151/pYEAS (VKPM B-7714)

VL2151/pYAHN (VKPM B-7748)

VL2151/pUC21 (VKPM B-7715)

5 W3350 *argE*::Tn10/pKA10, pYGGA (VKPM B-7716)

W3350 *argE*::Tn10/pKA10, pOK12 (VKPM B-7718)

10

15

20

25

30

35

40

45

50

55

SEQUENCE LISTING

5

10

<110> Ajinomoto Co., Inc.

15

<120> Method for Producing L-Amino Acid

<130> EPA-53281

<160> 24

20

<210> 1

<211> 27

<212> DNA

25

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer for amplifying Escherichia coli yahN gene

30

<400> 1

ggcgagctcc cagtaaccgg aaataag

27

35

<210> 2

<211> 27

<212> DNA

<213> Artificial Sequence

40

<220>

<223> Description of Artificial Sequence: primer for amplifying Escherichia coli yahN gene

45

<400> 2

cgctctagaa aggaccacgc attacgg

27

50

<210> 3

<211> 27

<212> DNA

55

<213> Artificial Sequence

5

<220>

<223> Description of Artificial Sequence: primer for amplifying Escherichia coli yeaS gene

10

<400> 3

ggcgagctca gattggttag catattc

27

15

<210> 4

<211> 27

<212> DNA

<213> Artificial Sequence

20

<220>

<223> Description of Artificial Sequence: primer for amplifying Escherichia coli yeaS gene

25

<400> 4

cggtctagaa tcagcgaaga atcaggg

27

30

<210> 5

<211> 27

<212> DNA

<213> Artificial Sequence

35

<220>

<223> Description of Artificial Sequence: primer for amplifying Escherichia coli yfiK gene

40

<400> 5

ggcgagctca tgttccgtgt cgggtac

27

45

<210> 6

<211> 27

<212> DNA

<213> Artificial Sequence

50

<220>

<223> Description of Artificial Sequence: primer for amplifying Escherichia coli yfiK gene

55

5 <400> 6
ggctctagat agcaagttac taagcgg 27

<210> 7
<211> 35
<212> DNA
10 <213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: primer for amplifying Escherichia
15 coli yggA gene

<400> 7
ctctgaattc tctcttatta gtttttctga ttgcc 35

20 <210> 8
<211> 38
<212> DNA
25 <213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: primer for amplifying Escherichia
30 coli yggA gene

<400> 8
cgtgacctgc agcgttctca cagcgcggta gcctttaa 38

35 <210> 9
<211> 672
<212> DNA
40 <213> Escherichia coli

<220>
<221> CDS
<222> (1)..(672)

45 <400> 9
atg atg cag tta gtt cac tta ttt atg gat gaa atc act atg gat cct 48
Met Met Gln Leu Val His Leu Phe Met Asp Glu Ile Thr Met Asp Pro
1 5 10 15
50 ttg cat gcc gtt tac ctg acc gta gga ctg ttc gtg att act ttt ttt 96
Leu His Ala Val Tyr Leu Thr Val Gly Leu Phe Val Ile Thr Phe Phe

55

EP 1 016 710 A2

| | | | | |
|----|---|-----|-----|-----|
| | 20 | 25 | 30 | |
| 5 | aat ccg gga gcc aat ctc ttt gtg gta gta caa acc agc ctg gct tcc | 144 | | |
| | Asn Pro Gly Ala Asn Leu Phe Val Val Val Gln Thr Ser Leu Ala Ser | | | |
| | 35 | 40 | 45 | |
| 10 | ggt cga cgc gca ggg gtg ctg acc ggg ctg ggc gtg gcg ctg ggc gat | 192 | | |
| | Gly Arg Arg Ala Gly Val Leu Thr Gly Leu Gly Val Ala Leu Gly Asp | | | |
| | 50 | 55 | 60 | |
| | gca ttt tat tcc ggg ttg ggt ttg ttt ggt ctt gca acg cta att acg | 240 | | |
| | Ala Phe Tyr Ser Gly Leu Gly Leu Phe Gly Leu Ala Thr Leu Ile Thr | | | |
| | 65 | 70 | 75 | 80 |
| 15 | cag tgt gag gag att ttt tcg ctt atc aga atc gtc ggc ggc gct tat | 288 | | |
| | Gln Cys Glu Glu Ile Phe Ser Leu Ile Arg Ile Val Gly Gly Ala Tyr | | | |
| | 85 | 90 | 95 | |
| 20 | ctc tta tgg ttt gcg tgg tgc agc atg cgc cgc cag tca aca ccg caa | 336 | | |
| | Leu Leu Trp Phe Ala Trp Cys Ser Met Arg Arg Gln Ser Thr Pro Gln | | | |
| | 100 | 105 | 110 | |
| | atg agc aca cta caa caa ccg att agc gcc ccc tgg tat gtc ttt ttt | 384 | | |
| | Met Ser Thr Leu Gln Gln Pro Ile Ser Ala Pro Trp Tyr Val Phe Phe | | | |
| | 115 | 120 | 125 | |
| 25 | cgc cgc gga tta att acc gat ctc tct aac ccg caa acc gtt tta ttt | 432 | | |
| | Arg Arg Gly Leu Ile Thr Asp Leu Ser Asn Pro Gln Thr Val Leu Phe | | | |
| | 130 | 135 | 140 | |
| 30 | ttt atc agt att ttc tca gta aca tta aat gcc gaa aca cca aca tgg | 480 | | |
| | Phe Ile Ser Ile Phe Ser Val Thr Leu Asn Ala Glu Thr Pro Thr Trp | | | |
| | 145 | 150 | 155 | 160 |
| | gca cgt tta atg gcc tgg gcg ggg att gtg ctc gca tca att atc tgg | 528 | | |
| | Ala Arg Leu Met Ala Trp Ala Gly Ile Val Leu Ala Ser Ile Ile Trp | | | |
| 35 | 165 | 170 | 175 | |
| | cga gtt ttt ctt agt cag gcg ttt tct ttg ccc gct gtg cgt cgt gct | 576 | | |
| | Arg Val Phe Leu Ser Gln Ala Phe Ser Leu Pro Ala Val Arg Arg Ala | | | |
| | 180 | 185 | 190 | |
| 40 | tat ggg cgt atg caa cgc gtt gcc agt cgg gtt att ggt gca att att | 624 | | |
| | Tyr Gly Arg Met Gln Arg Val Ala Ser Arg Val Ile Gly Ala Ile Ile | | | |
| | 195 | 200 | 205 | |
| 45 | ggt gta ttc gcg cta cgc ctg att tac gaa ggg gtg acg cag cgg tga | 672 | | |
| | Gly Val Phe Ala Leu Arg Leu Ile Tyr Glu Gly Val Thr Gln Arg | | | |
| | 210 | 215 | 220 | |

<210> 10

50 <211> 223

<212> PRT

<213> Escherichia coli

55

<400> 10
 Met Met Gln Leu Val His Leu Phe Met Asp Glu Ile Thr Met Asp Pro
 1 5 10 15
 Leu His Ala Val Tyr Leu Thr Val Gly Leu Phe Val Ile Thr Phe Phe
 20 25 30
 Asn Pro Gly Ala Asn Leu Phe Val Val Val Gln Thr Ser Leu Ala Ser
 35 40 45
 Gly Arg Arg Ala Gly Val Leu Thr Gly Leu Gly Val Ala Leu Gly Asp
 50 55 60
 Ala Phe Tyr Ser Gly Leu Gly Leu Phe Gly Leu Ala Thr Leu Ile Thr
 65 70 75 80
 Gln Cys Glu Glu Ile Phe Ser Leu Ile Arg Ile Val Gly Gly Ala Tyr
 85 90 95
 Leu Leu Trp Phe Ala Trp Cys Ser Met Arg Arg Gln Ser Thr Pro Gln
 100 105 110
 Met Ser Thr Leu Gln Gln Pro Ile Ser Ala Pro Trp Tyr Val Phe Phe
 115 120 125
 Arg Arg Gly Leu Ile Thr Asp Leu Ser Asn Pro Gln Thr Val Leu Phe
 130 135 140
 Phe Ile Ser Ile Phe Ser Val Thr Leu Asn Ala Glu Thr Pro Thr Trp
 145 150 155 160
 Ala Arg Leu Met Ala Trp Ala Gly Ile Val Leu Ala Ser Ile Ile Trp
 165 170 175
 Arg Val Phe Leu Ser Gln Ala Phe Ser Leu Pro Ala Val Arg Arg Ala
 180 185 190
 Tyr Gly Arg Met Gln Arg Val Ala Ser Arg Val Ile Gly Ala Ile Ile
 195 200 205
 Gly Val Phe Ala Leu Arg Leu Ile Tyr Glu Gly Val Thr Gln Arg
 210 215 220

 <210> 11
 <211> 639
 <212> DNA
 <213> Escherichia coli

 <220>
 <221> CDS
 <222> (1)..(639)

 <400> 11
 gtg ttc gct gaa tac ggg gtt ctg aat tac tgg acc tat ctg gtt ggg 48

EP 1 016 710 A2

| | | | | | | | | | | | | | | | | | |
|----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| | Met | Phe | Ala | Glu | Tyr | Gly | Val | Leu | Asn | Tyr | Trp | Thr | Tyr | Leu | Val | Gly | |
| | 1 | | | | 5 | | | | 10 | | | | | 15 | | | |
| 5 | gcc | att | ttt | att | gtg | ttg | gtg | cca | ggg | cca | aat | acc | ctg | ttt | gta | ctc | 96 |
| | Ala | Ile | Phe | Ile | Val | Leu | Val | Pro | Gly | Pro | Asn | Thr | Leu | Phe | Val | Leu | |
| | | | | 20 | | | | | 25 | | | | | 30 | | | |
| 10 | aaa | aat | agc | gtc | agt | agc | ggg | atg | aaa | ggc | ggg | tat | ctt | gcg | gcc | tgc | 144 |
| | Lys | Asn | Ser | Val | Ser | Ser | Gly | Met | Lys | Gly | Gly | Tyr | Leu | Ala | Ala | Cys | |
| | | | | 35 | | | | 40 | | | | | | 45 | | | |
| | ggg | gta | ttt | att | ggc | gat | gcg | gta | ttg | atg | ttt | ctg | gca | tgg | gct | gga | 192 |
| | Gly | Val | Phe | Ile | Gly | Asp | Ala | Val | Leu | Met | Phe | Leu | Ala | Trp | Ala | Gly | |
| 15 | | | | 50 | | | | 55 | | | | | 60 | | | | |
| | gtg | gcg | aca | tta | att | aag | acc | acc | ccg | ata | tta | ttc | aac | att | gta | cgt | 240 |
| | Val | Ala | Thr | Leu | Ile | Lys | Thr | Thr | Pro | Ile | Leu | Phe | Asn | Ile | Val | Arg | |
| | | | | 65 | | | | 70 | | | | | 75 | | | 80 | |
| 20 | tat | ctt | ggg | gcg | ttt | tat | ttg | ctc | tat | ctg | ggg | agt | aaa | att | ctt | tac | 288 |
| | Tyr | Leu | Gly | Ala | Phe | Tyr | Leu | Leu | Tyr | Leu | Gly | Ser | Lys | Ile | Leu | Tyr | |
| | | | | | 85 | | | | | 90 | | | | | 95 | | |
| | gcg | acc | ctg | aag | ggg | aaa | aat | agc | gag | gcc | aaa | tcc | gat | gag | ccc | caa | 336 |
| 25 | Ala | Thr | Leu | Lys | Gly | Lys | Asn | Ser | Glu | Ala | Lys | Ser | Asp | Glu | Pro | Gln | |
| | | | | 100 | | | | | 105 | | | | | 110 | | | |
| | tac | ggg | gct | att | ttt | aaa | cgc | gcg | tta | att | ttg | agc | ctg | act | aat | ccg | 384 |
| | Tyr | Gly | Ala | Ile | Phe | Lys | Arg | Ala | Leu | Ile | Leu | Ser | Leu | Thr | Asn | Pro | |
| 30 | | | | 115 | | | | 120 | | | | | 125 | | | | |
| | aaa | gcc | att | ttg | ttc | tat | gtg | tcg | ttt | ttc | gta | cag | ttt | atc | gat | gtt | 432 |
| | Lys | Ala | Ile | Leu | Phe | Tyr | Val | Ser | Phe | Phe | Val | Gln | Phe | Ile | Asp | Val | |
| | | | | 130 | | | | 135 | | | | 140 | | | | | |
| 35 | aat | gcc | cca | cat | acg | gga | att | tca | ttc | ttt | att | ctg | gcg | gcg | acg | ctg | 480 |
| | Asn | Ala | Pro | His | Thr | Gly | Ile | Ser | Phe | Phe | Ile | Leu | Ala | Ala | Thr | Leu | |
| | | | | | | 145 | | 150 | | | | 155 | | | 160 | | |
| | gaa | ctg | gtg | agt | ttc | tgc | tat | ttg | agc | ttc | ctg | att | ata | tct | ggg | gct | 528 |
| 40 | Glu | Leu | Val | Ser | Phe | Cys | Tyr | Leu | Ser | Phe | Leu | Ile | Ile | Ser | Gly | Ala | |
| | | | | | 165 | | | | | 170 | | | | 175 | | | |
| | ttt | gtc | acg | cag | tac | ata | cgt | acc | aaa | aag | aaa | ctg | gct | aaa | gtt | ggc | 576 |
| | Phe | Val | Thr | Gln | Tyr | Ile | Arg | Thr | Lys | Lys | Lys | Leu | Ala | Lys | Val | Gly | |
| 45 | | | | 180 | | | | | 185 | | | | | 190 | | | |
| | aac | tca | ctg | att | ggg | ttg | atg | ttc | gtg | ggg | ttc | gct | gcc | cga | ctg | gcg | 624 |
| | Asn | Ser | Leu | Ile | Gly | Leu | Met | Phe | Val | Gly | Phe | Ala | Ala | Arg | Leu | Ala | |
| | | | | 195 | | | | 200 | | | | | 205 | | | | |
| 50 | acg | ctg | caa | tcc | tga | | | | | | | | | | | | 639 |
| | Thr | Leu | Gln | Ser | | | | | | | | | | | | | |
| | | | | 210 | | | | | | | | | | | | | |

55

5 <210> 12
 <211> 212
 <212> PRT
 <213> Escherichia coli

 10 <400> 12
 Met Phe Ala Glu Tyr Gly Val Leu Asn Tyr Trp Thr Tyr Leu Val Gly
 1 5 10 15
 Ala Ile Phe Ile Val Leu Val Pro Gly Pro Asn Thr Leu Phe Val Leu
 20 25 30
 15 Lys Asn Ser Val Ser Ser Gly Met Lys Gly Gly Tyr Leu Ala Ala Cys
 35 40 45
 Gly Val Phe Ile Gly Asp Ala Val Leu Met Phe Leu Ala Trp Ala Gly
 50 55 60
 20 Val Ala Thr Leu Ile Lys Thr Thr Pro Ile Leu Phe Asn Ile Val Arg
 65 70 75 80
 Tyr Leu Gly Ala Phe Tyr Leu Leu Tyr Leu Gly Ser Lys Ile Leu Tyr
 85 90 95
 25 Ala Thr Leu Lys Gly Lys Asn Ser Glu Ala Lys Ser Asp Glu Pro Gln
 100 105 110
 Tyr Gly Ala Ile Phe Lys Arg Ala Leu Ile Leu Ser Leu Thr Asn Pro
 115 120 125
 30 Lys Ala Ile Leu Phe Tyr Val Ser Phe Phe Val Gln Phe Ile Asp Val
 130 135 140
 Asn Ala Pro His Thr Gly Ile Ser Phe Phe Ile Leu Ala Ala Thr Leu
 145 150 155 160
 35 Glu Leu Val Ser Phe Cys Tyr Leu Ser Phe Leu Ile Ile Ser Gly Ala
 165 170 175
 Phe Val Thr Gln Tyr Ile Arg Thr Lys Lys Lys Leu Ala Lys Val Gly
 180 185 190
 40 Asn Ser Leu Ile Gly Leu Met Phe Val Gly Phe Ala Ala Arg Leu Ala
 195 200 205
 Thr Leu Gln Ser
 210

45 <210> 13
 <211> 588
 <212> DNA
 <213> Escherichia coli

50 <220>
 <221> CDS

55

<222> (1)..(588)

5 <400> 13
 gtg aca ccg acc ctt tta agt gct ttt tgg act tac acc ctg att acc 48
 Met Thr Pro Thr Leu Leu Ser Ala Phe Trp Thr Tyr Thr Leu Ile Thr
 1 5 10 15
 10 gct atg acg cca gga ccg aac aat att ctc gcc ctt agc tct gct acg 96
 Ala Met Thr Pro Gly Pro Asn Asn Ile Leu Ala Leu Ser Ser Ala Thr
 20 25 30
 tcg cat gga ttt cgt caa agt acc cgc gtg ctg gca ggg atg agt ctg 144
 Ser His Gly Phe Arg Gln Ser Thr Arg Val Leu Ala Gly Met Ser Leu
 35 40 45
 gga ttt ttg att gtg atg tta ctg tgt gcg ggc att tca ttt tca ctg 192
 Gly Phe Leu Ile Val Met Leu Leu Cys Ala Gly Ile Ser Phe Ser Leu
 50 55 60
 20 gca gtg att gac ccg gca gcg gta cac ctt ttg agt tgg gcg ggg gcg 240
 Ala Val Ile Asp Pro Ala Ala Val His Leu Leu Ser Trp Ala Gly Ala
 65 70 75 80
 25 gca tat att gtc tgg ctg gcg tgg aaa atc gcc acc agc cca aca aag 288
 Ala Tyr Ile Val Trp Leu Ala Trp Lys Ile Ala Thr Ser Pro Thr Lys
 85 90 95
 gaa gac gga ctt cag gca aaa cca atc agc ttt tgg gcc agc ttt gct 336
 Glu Asp Gly Leu Gln Ala Lys Pro Ile Ser Phe Trp Ala Ser Phe Ala
 100 105 110
 30 ttg cag ttt gtg aac gtc aaa atc att ttg tac ggt gtt acg gca ctg 384
 Leu Gln Phe Val Asn Val Lys Ile Ile Leu Tyr Gly Val Thr Ala Leu
 115 120 125
 35 tcg acg ttt gtt ctg ccg caa aca cag gcg tta agc tgg gta gtt ggc 432
 Ser Thr Phe Val Leu Pro Gln Thr Gln Ala Leu Ser Trp Val Val Gly
 130 135 140
 gtc agc gtt ttg ctg gcg atg att ggg acg ttt ggc aat gtg tgc tgg 480
 Val Ser Val Leu Leu Ala Met Ile Gly Thr Phe Gly Asn Val Cys Trp
 145 150 155 160
 gcg ctg gcg ggg cat ctg ttt cag cga ttg ttt cgc cag tat ggt cgc 528
 Ala Leu Ala Gly His Leu Phe Gln Arg Leu Phe Arg Gln Tyr Gly Arg
 165 170 175
 45 cag tta aat atc gtg ctt gcc ctg ttg ctg gtc tat tgc gcg gta cgc 576
 Gln Leu Asn Ile Val Leu Ala Leu Leu Leu Val Tyr Cys Ala Val Arg
 180 185 190
 50 att ttc tat taa 588
 Ile Phe Tyr
 195

55

<210> 14
 <211> 195
 <212> PRT
 <213> Escherichia coli

<400> 14
 10 Met Thr Pro Thr Leu Leu Ser Ala Phe Trp Thr Tyr Thr Leu Ile Thr
 1 5 10 15
 Ala Met Thr Pro Gly Pro Asn Asn Ile Leu Ala Leu Ser Ser Ala Thr
 20 25 30
 15 Ser His Gly Phe Arg Gln Ser Thr Arg Val Leu Ala Gly Met Ser Leu
 35 40 45
 Gly Phe Leu Ile Val Met Leu Leu Cys Ala Gly Ile Ser Phe Ser Leu
 50 55 60
 20 Ala Val Ile Asp Pro Ala Ala Val His Leu Leu Ser Trp Ala Gly Ala
 65 70 75 80
 Ala Tyr Ile Val Trp Leu Ala Trp Lys Ile Ala Thr Ser Pro Thr Lys
 85 90 95
 25 Glu Asp Gly Leu Gln Ala Lys Pro Ile Ser Phe Trp Ala Ser Phe Ala
 100 105 110
 Leu Gln Phe Val Asn Val Lys Ile Ile Leu Tyr Gly Val Thr Ala Leu
 115 120 125
 30 Ser Thr Phe Val Leu Pro Gln Thr Gln Ala Leu Ser Trp Val Val Gly
 130 135 140
 Val Ser Val Leu Leu Ala Met Ile Gly Thr Phe Gly Asn Val Cys Trp
 145 150 155 160
 35 Ala Leu Ala Gly His Leu Phe Gln Arg Leu Phe Arg Gln Tyr Gly Arg
 165 170 175
 Gln Leu Asn Ile Val Leu Ala Leu Leu Leu Val Tyr Cys Ala Val Arg
 180 185 190
 40 Ile Phe Tyr
 195

<210> 15
 <211> 636
 <212> DNA
 <213> Escherichia coli

<220>
 <221> CDS
 <222> (1)..(636)

55

Leu Phe Ser
210

<210> 16

<211> 211

<212> PRT

<213> Escherichia coli

<400> 16

Met Phe Ser Tyr Tyr Phe Gln Gly Leu Ala Leu Gly Ala Ala Met Ile
1 5 10 15
Leu Pro Leu Gly Pro Gln Asn Ala Phe Val Met Asn Gln Gly Ile Arg
20 25 30
Arg Gln Tyr His Ile Met Ile Ala Leu Leu Cys Ala Ile Ser Asp Leu
35 40 45
Val Leu Ile Cys Ala Gly Ile Phe Gly Gly Ser Ala Leu Leu Met Gln
50 55 60
Ser Pro Trp Leu Leu Ala Leu Val Thr Trp Gly Gly Val Ala Phe Leu
65 70 75 80
Leu Trp Tyr Gly Phe Gly Ala Phe Lys Thr Ala Met Ser Ser Asn Ile
85 90 95
Glu Leu Ala Ser Ala Glu Val Met Lys Gln Gly Arg Trp Lys Ile Ile
100 105 110
Ala Thr Met Leu Ala Val Thr Trp Leu Asn Pro His Val Tyr Leu Asp
115 120 125
Thr Phe Val Val Leu Gly Ser Leu Gly Gly Gln Leu Asp Val Glu Pro
130 135 140
Lys Arg Trp Phe Ala Leu Gly Thr Ile Ser Ala Ser Phe Leu Trp Phe
145 150 155 160
Phe Gly Leu Ala Leu Leu Ala Ala Trp Leu Ala Pro Arg Leu Arg Thr
165 170 175
Ala Lys Ala Gln Arg Ile Ile Asn Leu Val Val Gly Cys Val Met Trp
180 185 190
Phe Ile Ala Leu Gln Leu Ala Arg Asp Gly Ile Ala His Ala Gln Ala
195 200 205
Leu Phe Ser
210

<210> 17

<211> 20

<212> DNA

<213> Artificial Sequence

5 <220>
 <223> Description of Artificial Sequence: primer for amplifying Escherichia
 coli yahN gene

 <400> 17
 gtgtggaacc gacgccgat 20
 10
 <210> 18
 <211> 23
 <212> DNA
 15 <213> Artificial Sequence

 <220>
 <223> Description of Artificial Sequence: primer for amplifying Escherichia
 20 coli yahN gene

 <400> 18
 tgttgtatgg tacggggttc gag 23
 25
 <210> 19
 <211> 20
 <212> DNA
 30 <213> Artificial Sequence

 <220>
 <223> Description of Artificial Sequence: primer for amplifying Escherichia
 35 coli yeaS gene

 <400> 19
 ctttgcaat cccgtctccc 20
 40
 <210> 20
 <211> 20
 <212> DNA
 45 <213> Artificial Sequence

 <220>
 <223> Description of Artificial Sequence: primer for amplifying Escherichia
 50 coli yeaS gene

 <400> 20
 55

gccccatgca taacggaaag 20

5 <210> 21
 <211> 26
 <212> DNA
 <213> Artificial Sequence

10 <220>
 <223> Description of Artificial Sequence: primer for amplifying Escherichia coli yfiK gene

15 <400> 21
 gaagatcttg taggccggat aaggcg 26

20 <210> 22
 <211> 20
 <212> DNA
 <213> Artificial Sequence

25 <220>
 <223> Description of Artificial Sequence: primer for amplifying Escherichia coli yfiK gene

30 <400> 22
 tggttttacc aattggccgc 20

35 <210> 23
 <211> 21
 <212> DNA
 <213> Artificial Sequence

40 <220>
 <223> Description of Artificial Sequence: primer for amplifying Escherichia coli yggA gene

45 <400> 23
 acttctcccg cgagccagtt c 21

50 <210> 24
 <211> 21
 <212> DNA
 <213> Artificial Sequence

55

<220>

<223> Description of Artificial Sequence: primer for amplifying *Escherichia coli* yggA gene

<400> 24

ggcaagctta ggcctctgt t

21

Claims

1. A bacterium belonging to the genus *Escherichia* and having an ability to produce an L-amino acid, wherein the ability to produce the L-amino acid is increased by increasing the amount of at least one protein produced in the bacterium and selected from the group consisting of the following proteins of (A) to (H):

- (A) a protein having an amino acid sequence shown in SEQ ID NO: 10 in Sequence Listing;
- (B) a protein which has an amino acid sequence including deletion, substitution, insertion, addition or inversion of one or several amino acids in the amino acid sequence shown in SEQ ID NO: 10 in Sequence Listing, and which has an activity of increasing the ability to produce the L-amino acid of the bacterium having the protein;
- (C) a protein having an amino acid sequence shown in SEQ ID NO: 12 in Sequence Listing;
- (D) a protein which has an amino acid sequence including deletion, substitution, insertion, addition or inversion of one or several amino acids in the amino acid sequence shown in SEQ ID NO: 12 in Sequence Listing, and which has an activity of increasing the ability to produce the L-amino acid of the bacterium having the protein;
- (E) a protein having an amino acid sequence shown in SEQ ID NO: 14 in Sequence Listing;
- (F) a protein which has an amino acid sequence including deletion, substitution, insertion, addition or inversion of one or several amino acids in the amino acid sequence shown in SEQ ID NO: 14 in Sequence Listing, and which has an activity of increasing the ability to produce the L-amino acid of the bacterium having the protein;
- (G) a protein having an amino acid sequence shown in SEQ ID NO: 16 in Sequence Listing; or
- (H) a protein which has an amino acid sequence including deletion, substitution, insertion, addition or inversion of one or several amino acids in the amino acid sequence shown in SEQ ID NO: 16 in Sequence Listing, and which has an activity of increasing the ability to produce the L-amino acid of the bacterium having the protein.

2. The bacterium according to claim 1, wherein said L-amino acid is L-lysine and the amount of at least one protein selected from the group consisting of said proteins (A) to (D), (G) and (H) is increased.
3. The bacterium according to claim 1, wherein said L-amino acid is L-glutamic acid.
4. The bacterium according to claim 1, wherein said L-amino acid is L-alanine and the amount of at least one protein selected from the group consisting of said proteins (C) and (D) is increased.
5. The bacterium according to claim 1, wherein said L-amino acid is L-valine and the amount of at least one protein selected from the group consisting of said proteins (C) and (D) is increased.
6. The bacterium according to claim 1, wherein said L-amino acid is L-histidine and the amount of at least one protein selected from the group consisting of said proteins (C) to (F) is increased.
7. The bacterium according to claim 1, wherein said L-amino acid is L-proline and the amount of at least one protein selected from the group consisting of said proteins (A) to (F) is increased.
8. The bacterium according to claim 1, wherein said L-amino acid is L-threonine and the amount of at least one protein selected from the group consisting of said proteins (E) and (F) is increased.

9. The bacterium according to claim 1, wherein said L-amino acid is L-arginine and the amount of at least one protein selected from the group consisting of said proteins (G) and (H) is increased.
- 5 10. The bacterium according to claim 1, wherein said L-amino acid is L-isoleucine and the amount of at least one protein selected from the group consisting of said proteins (C) and (D) is increased.
11. The bacterium according to any one of claims 1 to 10, wherein a copy number of a DNA coding for said protein in a cell is increased.
- 10 12. The bacterium according to claim 11, wherein said DNA is carried on a multicopy vector in the cell.
13. The bacterium according to claim 11, wherein said DNA is carried on a transposon in the cell.
14. A bacterium having the accession number VKPM B-7728, VKPM B-7729, VKPM B-7731, VKPM B-7730, VKPM B-7719, VKPM B-7722, VKPM B-7707, VKPM B-7712, VKPM B-7708, VKPM B-7753, VKPM B-7754, VKPM B-7752, VKPM B-7713, VKPM B-7714, VKPM B-7748, VKPM B-7715, VKPM B-7716 or VKPM B-7718.
- 15 15. A method for producing an L-amino acid, comprising the steps of:
- 20 cultivating the bacterium as defined in any of the claims 1 to 14, in a culture medium, to produce and accumulate the L-amino acid in the medium, and recovering the L-amino acid from the medium.

25

30


35

40

45

50

55



Europäisches Patentamt
European Patent Office
Office européen des brevets



(11) **EP 1 016 710 A3**

(12) **EUROPEAN PATENT APPLICATION**

(88) Date of publication A3:
06.09.2000 Bulletin 2000/36

(51) Int. Cl.⁷: **C12N 1/20, C12N 15/11,
C07K 14/245**

(43) Date of publication A2:
05.07.2000 Bulletin 2000/27

(21) Application number: **99125263.6**

(22) Date of filing: **17.12.1999**

(84) Designated Contracting States:
**AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE**
Designated Extension States:
AL LT LV MK RO SI

(30) Priority: **30.12.1998 RU 98124016
09.03.1999 RU 99104431**

(71) Applicant: **Ajinomoto Co., Inc.
Tokyo (JP)**

(72) Inventors:
• **Livshits, Vitaliy Arkadievich
Moscow, 113208 (RU)**

- **Zakataeva, Natalia Pavlovna
Moscow, 117421 (RU)**
- **Nakanishi, Kazuo
Sakae-ku, Yokohama-shi, 247-0014 (JP)**
- **Aleshin, Vladimir Veniaminovich
Borovsk, 249010 (RU)**
- **Troshin, Petr Vladimirovich
Moscow, 115561 (RU)**
- **Tokhmakova, Irina Lyvovna
143513 Moscow region (RU)**

(74) Representative:
**Strehl Schübel-Hopf & Partner
Maximilianstrasse 54
80538 München (DE)**

(54) **Method for producing L-amino acids**

(57) A bacterium belonging to the genus *Escherichia* and having an ability to produce an L-amino acid, wherein the ability to produce the L-amino acid is increased by increasing an expression amount of an L-amino acid excretion protein, and a method for producing the L-amino acid using the bacterium.

EP 1 016 710 A3



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number
EP 99 12 5263

| DOCUMENTS CONSIDERED TO BE RELEVANT | | | |
|---|--|---|---|
| Category | Citation of document with indication, where appropriate, of relevant passages | Relevant to claim | CLASSIFICATION OF THE APPLICATION (Int.Cl.7) |
| Y | ZAKATAEVA, N. P. ET AL: "Characterization of a pleiotropic mutation that confers upon Escherichia coli cells resistance to high concentrations of homoserine and threonine." FASEB JOURNAL, (1997) VOL. 11, NO. 9, PP. A935. MEETING INFO.: 17TH INTERNATIONAL CONGRESS OF BIOCHEMISTRY AND MOLECULAR BIOLOGY IN CONJUNCTION WITH THE ANNUAL MEETING OF THE AMERICAN SOCIETY FOR BIOCHEMISTRY AND MOLECULAR BIOLOGY SAN FRANCISCO, CALI, XP002136826 | 1-8, 10-15 | C12N1/20 C12N15/11 C07K14/245 |
| A | "Y" for inventions 1-3; "A" for invention 4. * the whole document * | 9 | |
| Y | DATABASE EMBL/SWISSPROT [Online] ID: YAHN_ECOLI, 1 November 1997 (1997-11-01) DUNCAN ET AL.: "YAHN" XP002136827 Invention 1. See annotation - "Similarity: Belongs to the RHT family." * abstract * | 1-3,7, 11-15 | TECHNICAL FIELDS SEARCHED (Int.Cl.7) C12N |
| T | EP 0 994 190 A (AJINOMOTO KK) 19 April 2000 (2000-04-19) All inventions. * the whole document * | 1-15 | |
| Y | DATABASE EMBL/SWISSPROT [Online] ID: YEAS_ECOLI, 15 July 1998 (1998-07-15) ITO ET AL.: "YEAS" XP002142361 Invention 2. See annotation - "Similarity: Belongs to the RHT family." * abstract * | 1-7, 10-15 | |
| The present search report has been drawn up for all claims | | | |
| Place of search MUNICH | | Date of completion of the search 12 July 2000 | Examiner Sprinks, M |
| CATEGORY OF CITED DOCUMENTS X: particularly relevant if taken alone Y: particularly relevant if combined with another document of the same category A: technological background O: non-written disclosure P: intermediate document | | T: theory or principle underlying the invention E: earlier patent document, but published on, or after the filing date D: document cited in the application L: document cited for other reasons &: member of the same patent family, corresponding document | |

EPO FORM 1503 03.02 (P04021)



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number
EP 99 12 5263

| DOCUMENTS CONSIDERED TO BE RELEVANT | | | |
|--|---|---|--|
| Category | Citation of document with indication, where appropriate, of relevant passages | Relevant to claim | CLASSIFICATION OF THE APPLICATION (Int.Cl.7) |
| Y | <p>DATABASE EMBL/SWISSPROT [Online] ID: YFIK_ECOLI, 1 October 1994 (1994-10-01) NASHIMOTO: "YFIK" XP002142362 Invention 3. See annotation - "Similarity: Belongs to the RHT family." * abstract *</p> | 1,3,6-8, 11-15 | |
| A | <p>DATABASE EMBL/SWISSPROT [Online] ID: YGGA_ECOLI, 1 October 1989 (1989-10-01) ALEFOUNDER ET AL.: "YGGA" XP002142363 Invention 4. See annotation - "Similarity: Belongs to the LYSE/YGGA family." * abstract *</p> | 1-3,9, 11-15 | |
| | | | TECHNICAL FIELDS SEARCHED (Int.Cl.7) |
| The present search report has been drawn up for all claims | | | |
| Place of search MUNICH | | Date of completion of the search 12 July 2000 | Examiner Sprinks, M |
| <p>CATEGORY OF CITED DOCUMENTS</p> <p>X: particularly relevant if taken alone Y: particularly relevant if combined with another document of the same category A: technological background O: non-written disclosure P: intermediate document</p> | | <p>T: theory or principle underlying the invention E: earlier patent document, but published on, or after the filing date D: document cited in the application L: document cited for other reasons</p> <p>&: member of the same patent family, corresponding document</p> | |

EPO FORM 1503 03/02 (P04C01)

**ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO. .**

EP 99 12 5263

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report. The members are as contained in the European Patent Office EDP file on The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

12-07-2000

| Patent document cited in search report | Publication date | Patent family member(s) | Publication date |
|---|---------------------|---------------------------------|--------------------------|
| EP 0994190 A | 19-04-2000 | AU 4755099 A JP 2000116390 A | 20-04-2000 25-04-2000 |
| ----- | | | |

EPO FORM P0459

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82

THIS PAGE BLANK (USPTO)